

THE γ -POSITIVITY OF BASIC EULERIAN POLYNOMIALS VIA GROUP ACTIONS

ZHICONG LIN AND JIANG ZENG

ABSTRACT. We provide combinatorial interpretation for the γ -coefficients of the basic Eulerian polynomials that enumerate permutations by the excedance statistic and the major index as well as the corresponding γ -coefficients for derangements. Our results refine the classical γ -positivity results for the Eulerian polynomials and the derangement polynomials. The main tools are Brändén's modified Foata–Strehl action on permutations and the recent triple statistic $(\text{des}, \text{rix}, \text{aid})$ equidistributed with $(\text{exc}, \text{fix}, \text{maj})$.

1. INTRODUCTION

Any polynomial $h(t) = \sum_{i=0}^n h_i t^i$ with symmetric coefficient sequence ($h_{n-i} = h_i$) can be expanded uniquely as $h(t) = \sum_{k=0}^{\lfloor n/2 \rfloor} \gamma_k t^k (1+t)^{n-2k}$, and is said to be γ -positive if $\gamma_k \geq 0$. An interesting problem is then to see whether there are any combinatorial or geometric meaning of the γ -coefficients when they are positive, see [2, 5, 12, 13]. It is well known that Eulerian polynomials are γ -positive, see (1.6) below. In this work we shall study two generalizations of the γ -positivity of Eulerian polynomials.

A permutation σ of $[n] := \{1, 2, \dots, n\}$ is a bijection from $[n]$ to $[n]$, which will be identified with the word $\sigma = \sigma_1 \cdots \sigma_n$, where $\sigma_i = \sigma(i)$. We denote by $\mathfrak{S}_n$ the set of all permutations of $[n]$. Given a $\sigma \in \mathfrak{S}_n$, an integer $i \in [n]$ is an *excedance* (resp. *fixed point*) of σ if $\sigma_i > i$ (resp. $\sigma_i = i$). Denote by $\text{exc}(\sigma)$ and $\text{fix}(\sigma)$ the number of excedances and fixed points of σ , respectively. Let $\text{maj}(\sigma) := \sum_{\sigma_i > \sigma_{i+1}} i$ be the major index of σ . Shareshian and Wachs [15, 16] introduced and studied the *basic Eulerian polynomials*

$$A_n(t, r, q) := \sum_{\sigma \in \mathfrak{S}_n} t^{\text{exc}(\sigma)} r^{\text{fix}(\sigma)} q^{\text{maj}(\sigma) - \text{exc}(\sigma)}.$$

In [16, Remark 5.5] they noticed that using an unpublished result of Gessel they could prove, among other things, that

(1) there are polynomials $\gamma_{n,k}(q)$ in $\mathbb{N}[q]$ such that

$$A_n(t, 1, q) = \sum_{k=0}^{\lfloor n/2 \rfloor} \gamma_{n,k}(q) t^k (1+t)^{n-2k}. \quad (1.1)$$

Date: May 1, 2015.

Key words and phrases. Eulerian polynomials; γ -positivity; derangements; major index; inversions; admissible inversions.

(2) there are polynomials $\tilde{\gamma}_{n,k}(q)$ in $\mathbb{N}[q]$ such that

$$A_n(t, 0, q) = \sum_{k=0}^{\lfloor (n-1)/2 \rfloor} \tilde{\gamma}_{n,k}(q) t^k (1+t)^{n-1-2k}. \quad (1.2)$$

They have also two refined versions of (1.2), that we shall discuss in the last section, see (6.4) and (6.5).

The aim of this paper is to provide combinatorial interpretations for the coefficients $\gamma_{n,k}(q)$ and $\tilde{\gamma}_{n,k}(q)$. Several q -analogs of Eulerian polynomials with combinatorial meanings have been studied in the literature (see [15] and references therein) and various extensions of (1.6) and (1.8) have already been obtained in [2, 9, 17, 18]. We first review some partial and related recent results on this topic.

Let $\sigma = \sigma_1 \sigma_2 \dots \sigma_n$ be permutation in $\mathfrak{S}_n$ and by convention $\sigma_0 = \sigma_{n+1} = +\infty$. We call σ_i ($1 \leq i \leq n$) a *double descent* (resp. *double ascent*, *peak*, *valley*) of σ if $\sigma_{i-1} > \sigma_i > \sigma_{i+1}$ (resp. $\sigma_{i-1} < \sigma_i < \sigma_{i+1}$, $\sigma_{i-1} < \sigma_i > \sigma_{i+1}$, $\sigma_{i-1} > \sigma_i < \sigma_{i+1}$). Denote by $\text{dd}(\sigma)$ (resp. $\text{da}(\sigma)$, $\text{peak}(\sigma)$, $\text{valley}(\sigma)$) the number of double descents (resp. double ascents, peaks, valleys) of σ . For each $\sigma \in \mathfrak{S}_n$ let $\text{cda}(\sigma) := |\{i : \sigma^{-1}(i) < i < \sigma(i)\}|$ be the number of *double excedances* or *cyclic double ascents* of σ and $\text{cyc}(\sigma)$ denote the number of cycles of σ . For $1 \leq k \leq n$ define the sets

$$\begin{aligned} \mathfrak{D}_{n,k} &:= \{\sigma \in \mathfrak{S}_n : \text{dd}(\sigma) = 0, \text{des}(\sigma) = k\}, \\ \tilde{\mathfrak{D}}_{n,k} &:= \{\sigma \in \mathfrak{D}_{n,k-1} : \sigma_{n-1} < \sigma_n\}, \\ \mathcal{E}_{n,k} &:= \{\sigma \in \mathfrak{S}_n : \text{fix}(\sigma) = 0, \text{cda}(\sigma) = 0, \text{exc}(\sigma) = k\}. \end{aligned}$$

For example, for $n = 4$, we have

$$\mathfrak{D}_{4,0} = \{1234\} \quad \text{and} \quad \mathfrak{D}_{4,1} = \{1324, 1423, 2314, 2413, 3412, 1243, 1342, 2341\}; \quad (1.3)$$

$$\tilde{\mathfrak{D}}_{4,1} = \{1234\} \quad \text{and} \quad \tilde{\mathfrak{D}}_{4,2} = \{1324, 1423, 2314, 2413, 3412\}; \quad (1.4)$$

$$\mathcal{E}_{4,1} = \{4123\} \quad \text{and} \quad \mathcal{E}_{4,2} = \{2143, 2413, 3412, 4312, 4321\}. \quad (1.5)$$

A classical result of Foata and Schützenberger [5] states that the Eulerian polynomials have the following γ -expansion, which implies both the *symmetry* and *unimodality* (see for instance [16] for definitions) of the sequence of Eulerian numbers.

Theorem 1 (Foata–Schützenberger). *One has*

$$A_n(t, 1, 1) = \sum_{k=0}^{\lfloor (n-1)/2 \rfloor} |\mathfrak{D}_{n,k}| t^k (1+t)^{n-1-2k}. \quad (1.6)$$

Shin and Zeng [17, Theorem 11] have recently proved the following result via continued fractions.

Theorem 2 (Shin–Zeng). *One has*

$$\sum_{\substack{\sigma \in \mathfrak{S}_n \\ \text{fix}(\sigma)=0}} \beta^{\text{cyc}(\sigma)} t^{\text{exc}(\sigma)} = \sum_{k=1}^{\lfloor n/2 \rfloor} \left(\sum_{\sigma \in \mathcal{E}_{n,k}} \beta^{\text{cyc}(\sigma)} \right) t^k (1+t)^{n-2k}. \quad (1.7)$$

In particular, we have the derangement analogue of (1.6)

$$A_n(t, 0, 1) = \sum_{k=1}^{\lfloor n/2 \rfloor} |\mathcal{E}_{n,k}| t^k (1+t)^{n-2k}. \quad (1.8)$$

For $n = 4$, it follows from (1.5) that

$$\begin{aligned} \sum_{\substack{\sigma \in \mathfrak{S}_n \\ \text{fix}(\sigma)=0}} \beta^{\text{cyc}(\sigma)} t^{\text{exc}(\sigma)} &= \beta t(1+t)^2 + (2\beta + 3\beta^2) t^2 \\ &= \beta t + 4\beta t^2 + 3\beta^2 t^2 + \beta t^3. \end{aligned}$$

A permutation $\sigma \in \mathfrak{S}_n$ is called *alternating* if $\sigma_1 < \sigma_2 > \sigma_3 < \sigma_4 > \dots$. An *inversion* of a permutation $\sigma \in \mathfrak{S}_n$ is a pair (σ_i, σ_j) such that $1 \leq i < j \leq n$ and $\sigma_i > \sigma_j$. Let $\text{inv}(\sigma)$ be the number of inversions of σ . The set of alternating permutations of order n is denoted by $\mathfrak{A}_n$. When $t = -1$ Foata and Han [4, Theorem 1] proved the following q -analogue of (1.6) and (1.8).

Theorem 3 (Foata-Han). *One has $A_{2n}(-1, 1, q) = A_{2n-1}(-1, 0, q) = 0$ ($n \geq 1$) and*

$$\begin{aligned} A_{2n+1}(-1, 1, q) &= (-1)^n \sum_{\sigma \in \mathfrak{A}_{2n+1}} q^{\text{inv}(\sigma)} \quad (n \geq 0); \\ A_{2n}(-1, 0, q) &= (-1)^n \sum_{\sigma \in \mathfrak{A}_{2n}} q^{\text{inv}(\sigma)} \quad (n \geq 0). \end{aligned}$$

Our main results, Theorems 4 and 5, are q -analogues of (1.6) and (1.8) for general t .

Theorem 4. *We have*

$$A_n(t, 1, q) = \sum_{k=0}^{\lfloor (n-1)/2 \rfloor} \left(\sum_{\sigma \in \mathfrak{D}_{n,k}} q^{\text{inv}(\sigma)} \right) t^k (1+t)^{n-1-2k}. \quad (1.9)$$

For $n = 4$, it follows from (1.3) that

$$\begin{aligned} A_4(t, 1, q) &= (1+t)^3 + (2q + 3q^2 + 2q^3 + q^4)t(1+t) \\ &= 1 + (3 + 2q + 3q^2 + 2q^3 + q^4)t + (3 + 2q + 3q^2 + 2q^3 + q^4)t^2 + t^3. \end{aligned}$$

Theorem 5. *We have*

$$A_n(t, 0, q) = \sum_{k=1}^{\lfloor n/2 \rfloor} \left(\sum_{\sigma \in \mathfrak{D}_{n,k}} q^{\text{inv}(\sigma)} \right) t^k (1+t)^{n-2k}. \quad (1.10)$$

For $n = 4$, it follows from (1.4) that

$$\begin{aligned} A_4(t, 0, q) &= t(1+t)^2 + (q + 2q^2 + q^3 + q^4)t^2 \\ &= t + (2 + q + 2q^2 + q^3 + q^4)t^2 + t^3. \end{aligned}$$

Obviously Theorem 4 is a q -analogue of Theorem 1. To show that (1.10) reduces to (1.8), we will give a bijection from $\widetilde{\mathfrak{D}}_{n,k}$ to $\mathcal{E}_{n,k}$ in Remark 3. Since $\mathfrak{A}_{2n+1} = \mathfrak{D}_{2n+1,n}$ and $\mathfrak{A}_{2n} = \widetilde{\mathfrak{D}}_{2n,n}$, Theorems 4 and 5 with $t = -1$ reduce immediately to Theorem 3.

Remark 1. For any $\sigma \in \mathfrak{S}_n$ let $\text{DES}(\sigma) = \{i \in [n-1] : \sigma_i > \sigma_{i+1}\}$ be its descent set. According to a result of Foata and Schützenberger [6], for any $S \subseteq [n-1]$,

$$\sum q^{\text{inv}(\sigma)} = \sum q^{\text{imaj}(\sigma)} \quad (\sigma \in \mathfrak{S}_n \text{ and } \text{DES}(\sigma) = S),$$

where $\text{imaj}(\sigma) = \text{maj}(\sigma^{-1})$. Hence we can replace the statistic inv by imaj in Theorems 4 and 5.

We will prove Theorems 4 and 5 by making use of Brändén's modified Foata–Strehl action, [2, 7] (see also [14]), and an alternative interpretation of the basic Eulerian polynomials involving the descent statistic, that we recall below.

Definition 1. Let $w = w_1 w_2 \cdots w_k$ be a word of length k with distinct letters from $[n]$. An admissible inversion of w is a pair (w_i, w_j) such that $1 \leq i < j \leq k$ and $w_i > w_j$ and satisfies either of the following conditions:

- $1 < i$ and $w_{i-1} < w_i$ or
- there is some l such that $i < l < j$ and $w_i < w_l$.

Let $\text{ai}(w)$ be the number of admissible inversions of w . The statistic “rix” is defined recursively as follows: $\text{rix}(\emptyset) = 0$, if $w_i = \max\{w_1, w_2, \dots, w_k\}$ ($1 \leq i \leq k$), then

$$\text{rix}(w) := \begin{cases} 0, & \text{if } i = 1 < k; \\ 1 + \text{rix}(w_1 w_2 \cdots w_{k-1}), & \text{if } i = k; \\ \text{rix}(w_{i+1} w_{i+2} \cdots w_k), & \text{if } 1 < i < k. \end{cases}$$

Note that $\text{rix}(w_1) = 1$. If $w = 291753468$, then there are 14 inversions, except $(5, 3)$ and $(5, 4)$, all others are admissible, hence $\text{ai}(w) = 12$; for the computation of $\text{rix}(w)$, by looking for the greatest letters in the words successively we have

$$\text{rix}(w) = \text{rix}(1753468) = 1 + \text{rix}(175346) = 1 + \text{rix}(5346) = 2 + \text{rix}(534) = 2.$$

We will need the following interpretation of the basic Eulerian polynomials [10, Theorem 8] (see also [3] for an equivalent version), of which the special $r = 1$ case was first proved in [11] using the Rees product of posets.

Lemma 6 ([3, 10, 16]). One has

$$A_n(t, r, q) = \sum_{\sigma \in \mathfrak{S}_n} t^{\text{des}(\sigma)} r^{\text{rix}(\sigma)} q^{\text{ai}(\sigma)}. \quad (1.11)$$




FIGURE 1. MFS-actions on 65137428

The rest of this paper is organized as follows. In Section 2, after recalling the Modified Foata–Strehl action, we prove Theorem 4. In Section 3, we introduce an alternative algorithm to compute the statistic rix , which connects another model $\mathcal{R}_{n,k}^0$ (see (3.2)) to $\mathcal{E}_{n,k}$. Proof of Theorem 5 will be given in Section 4. We derive the recurrence relations and the generating functions of the γ -coefficients in (1.9) and (1.10) in Section 5. We conclude the paper with some further remarks in Section 6.

2. PROOF OF THEOREM 4

Let $\sigma \in \mathfrak{S}_n$, for any $x \in [n]$, the x -factorization of σ reads $\sigma = w_1 w_2 x w_3 w_4$, where w_2 (resp. w_3) is the maximal contiguous subword immediately to the left (resp. right) of x whose letters are all smaller than x . Following Foata and Strehl [7] we define the action φ_x by

$$\varphi_x(\sigma) = w_1 w_3 x w_2 w_4.$$

For instance, if $x = 4$ and $\sigma = 2743156 \in \mathfrak{S}_7$, then $w_1 = 27$, $w_2 = \emptyset$, $w_3 = 31$ and $w_4 = 56$. Thus $\varphi_x(\sigma) = 2731456$. Clearly, φ_x is an involution acting on $\mathfrak{S}_n$ and it is not hard to see that φ_x and φ_y commute for all $x, y \in [n]$. Brändén [2] modified φ_x to be

$$\varphi'_x(\sigma) := \begin{cases} \varphi_x(\sigma), & \text{if } x \text{ is a double ascent or double descent of } \sigma; \\ \sigma, & \text{if } x \text{ is a valley or a peak of } \sigma. \end{cases}$$

Again it is clear that φ'_x 's are involutions and commute. For any subset $S \subseteq [n]$ we can then define the function $\varphi'_S : \mathfrak{S}_n \rightarrow \mathfrak{S}_n$ by

$$\varphi'_S(\sigma) = \prod_{x \in S} \varphi'_x(\sigma).$$

Hence the group $\mathbb{Z}_2^n$ acts on $\mathfrak{S}_n$ via the functions φ'_S , $S \subseteq [n]$. This action will be called the *Modified Foata–Strehl action* (MFS-action for short) as depicted in Fig. 1. We first show that the statistic “ai” is invariant under the MFS-action.

Lemma 7. *Let $\sigma \in \mathfrak{S}_n$. For each $x \in [n]$, we have $\text{ai}(\sigma) = \text{ai}(\varphi'_x(\sigma))$.*

Proof. If x is a peak or a valley of σ , then $\varphi'_x(\sigma) = \sigma$ and the result is true. If x is a double descent of σ , then w_2 is empty in the x -factorization of $\sigma = w_1 x w_3 w_4$ and there is

no admissible inversions of σ between x and the word w_3 . As $\varphi'_x(\sigma) = w_1 w_3 x w_4$, there is no inversions of $\varphi'_x(\sigma)$ between the word w_3 and x . Let $(\sigma_i, \sigma_j) \notin \{(x, y) : y \text{ is a letter in } w_3\}$ be a pair of σ such that $i < j$. We claim that (σ_i, σ_j) is an admissible inversion of σ if and only if it is an admissible inversion of $\varphi'_x(\sigma)$, from which the result follows.

For a word w , we write $a \in w$ if a is a letter in w . To check the claim, there are 6 cases to be considered: (1) $\sigma_i \in w_1$ and $\sigma_j \in w_1$; (2) $\sigma_i \in w_1$ and $\sigma_j \in x w_3$; (3) $\sigma_i \in w_1$ and $\sigma_j \in w_4$; (4) $\sigma_i \in w_3$ and $\sigma_j \in w_3$; (5) $\sigma_i \in x w_3$ and $\sigma_j \in w_4$; (6) $\sigma_i \in w_4$ and $\sigma_j \in w_4$. We will only show case (2), other cases are similar. If (σ_i, σ_j) is an admissible inversion of σ , then $\sigma_{i-1} < \sigma_i > \sigma_j$ or $\sigma_j < \sigma_i < \sigma_k$ for some $i < k < j$. Clearly, (σ_i, σ_j) is an admissible of $\varphi'_x(\sigma)$ if $\sigma_k \neq x$. Otherwise if $\sigma_k = x$, then we denote x' the last letter of w_1 and consider the triple (σ_i, x', σ_j) . This indicates that (σ_i, σ_j) is an admissible inversion of $\varphi'_x(\sigma)$, since $x' > x > \sigma_i$. To show that, if (σ_i, σ_j) is an admissible inversion of $\varphi'(\sigma)$ then (σ_i, σ_j) is an admissible inversion of σ , is similar and we omit. This finishes the proof of our claim in case (2). $\square$

For any permutation $\sigma \in \mathfrak{S}_n$, let $\text{Orb}(\sigma) = \{g(\sigma) : g \in \mathbb{Z}_2^n\}$ be the orbit of σ under the MFS-action. The MFS-action divides the set $\mathfrak{S}_n$ into disjoint orbits. Moreover, for $\sigma \in \mathfrak{S}_n$, if x is a double descent of σ , then x is a double ascent of $\varphi'_x(\sigma)$. Hence, there is a unique permutation in each orbit which has no double descent. Now, let $\bar{\sigma}$ be such a unique element in $\text{Orb}(\sigma)$, then $\text{da}(\bar{\sigma}) = n - \text{peak}(\bar{\sigma}) - \text{valley}(\bar{\sigma})$ and $\text{des}(\bar{\sigma}) = \text{peak}(\bar{\sigma}) = \text{valley}(\bar{\sigma}) - 1$. Thus

$$\sum_{\pi \in \text{Orb}(\sigma)} q^{\text{ai}(\pi)} t^{\text{des}(\pi)} = q^{\text{ai}(\bar{\sigma})} t^{\text{des}(\bar{\sigma})} (1+t)^{\text{da}(\bar{\sigma})} = q^{\text{ai}(\bar{\sigma})} t^{\text{des}(\bar{\sigma})} (1+t)^{n-1-2\text{des}(\bar{\sigma})}.$$

Therefore, by (1.11), we have

$$A_n(t, 1, q) = \sum_{\sigma \in \mathfrak{S}_n} t^{\text{des}(\sigma)} q^{\text{ai}(\sigma)} = \sum_{k=0}^{\lfloor (n-1)/2 \rfloor} \left(\sum_{\sigma \in \mathfrak{D}_{n,k}} p^{\text{ai}(\sigma)} \right) t^k (1+t)^{n-1-2k}.$$

Thus Theorem 4 is a consequence of the following result.

Lemma 8. *For each $\sigma \in \mathfrak{D}_{n,k}$, we have $\text{ai}(\sigma) = \text{inv}(\sigma)$.*

Proof. Let $\sigma \in \mathfrak{S}_n$ be a permutation without double descent. Let (σ_i, σ_j) be an inversion of σ . If $i = 1$, then $\sigma_1 < \sigma_2$ and so (σ_i, σ_j) is an admissible inversion of σ . Assume that $i \geq 2$. If $\sigma_{i-1} < \sigma_i$ then (σ_i, σ_j) is an admissible inversion of σ ; if $\sigma_{i-1} > \sigma_i$, then we must have $\sigma_{i+1} > \sigma_i$, otherwise σ_i will be a double descent of σ , hence (σ_i, σ_j) is also an admissible inversion of σ . This shows that $\text{ai}(\sigma) \geq \text{inv}(\sigma)$. As $\text{ai}(\sigma) \leq \text{inv}(\sigma)$ by definition, we are done. $\square$

3. RIX-FACTORIZATION

We first give a slightly less recursive description of the “rix” statistic. Let $w = w_1 \cdots w_k$ be a word with distinct letters over the alphabet $[n]$. We say that w_i is a *descent top* of

w if $w_i > w_{i+1}$. If $k = 1$ the word w is called a L-hook, if $k \geq 2$ the word w is called a L-hook (resp. F-hook) if the last (resp. first) letter w_k (resp. w_1) is the greatest letter of w .

Definition 2 (Rix-factorization). *Each permutation $\sigma \in \mathfrak{S}_n$ can be factorized as*

$$\sigma = \alpha_1 \alpha_2 \cdots \alpha_i \beta, \quad (3.1)$$

where each α_j ($1 \leq j \leq i$) is a L-hook of length ≥ 2 (possibly empty) and β is a L-hook or F-hook, by applying the following algorithm:

- (i) $w \leftarrow \sigma$; $i \leftarrow 0$;
- (ii) if w is an increasing word, let $\beta = w$ and we get (3.1); otherwise, $i \leftarrow i + 1$, let x be the greatest descent top of w and write $w = w' x w''$ for some subwords w', w'' ;
- (iii) if $w' = \emptyset$, let $\beta = w$ and we get (3.1); otherwise, let $\alpha_i = w' x$, $w \leftarrow w''$ and go to (ii).

The factorization (3.1) will be called the rix-factorization of σ . Denote by $\beta_1(\sigma)$ the first letter of β . Then a letter in the maximal increasing suffix of β that is not smaller than $\beta_1(\sigma)$ will be called a rixed point of σ . Denote by $\text{RIX}(\sigma)$ the set of all rixed points of σ .

For example, the algorithm gives

$$\begin{aligned} \sigma &:= 21879354610 = 21879|35|\beta \quad \text{with} \quad \beta = 4610 \text{ (L-hook)}, \\ \tau &:= 61108497253 = 6110|849|\beta \quad \text{with} \quad \beta = 7253 \text{ (F-hook)}, \\ \pi &:= 11047625389 = 110|47|\beta \quad \text{with} \quad \beta = 625389 \text{ (L-hook)}. \end{aligned}$$

We have

$$\begin{aligned} \beta_1(\sigma) &= 4, \quad \text{RIX}(\sigma) = \{4, 6, 10\}; \\ \beta_1(\tau) &= 7, \quad \text{RIX}(\tau) = \emptyset; \\ \beta_1(\pi) &= 6, \quad \text{RIX}(\pi) = \{8, 9\}. \end{aligned}$$

Proposition 9. *Let $\sigma \in \mathfrak{S}_n$ with rix-factorization (3.1). If x_k is the last letter of α_k ($1 \leq k \leq i$), then*

- (i) $x_1 > x_2 > \cdots > x_i > \beta_1(\sigma)$.
- (ii) $\text{rix}(\sigma) = 0$ if and only if β is an F-hook.
- (iii) $\text{rix}(\sigma) = |\text{RIX}(\sigma)|$.

Proof. (i) For $k = 1, \dots, i$, since x_k is the greatest descent top of $\alpha_k \cdots \alpha_i \beta$, it is obvious that $x_k > x_{k+1}$, where $x_{i+1} = \beta_1(\sigma)$. (ii) If β is an F-hook (so the length of β is at least 2), when applying the algorithmic definition to σ for computing the statistic “rix”, the successive greatest letters are $x_1, x_2, \dots, x_i$ and $\beta_1(\sigma)$, so $\text{rix}(\sigma) = 0$ because $\beta_1(\sigma)$ is the first letter of β . The “only if” part will follow from (iii). Indeed, if β is an L-hook, when applying the algorithmic definition to σ for computing the statistic “rix”, the greatest letters are the letters in

$$\text{RIX}(\sigma) \cup \{x_1, x_2, \dots, x_i, \beta_1(\sigma)\},$$

read in decreasing order, but only the letters in $\text{RIX}(\sigma)$ will contribute to the statistic “rix”. $\square$

The rix-factorization is reminiscent of Gessel’s hook factorization [8], which decomposes each permutation into pieces of hooks, uniquely. Indeed we have the following result.

Proposition 10. *The rix-factorization (3.1) is unique provided condition (i) in Proposition 9 holds and $\beta_1(\sigma)$ is the greatest descent top of β (whenever β is not an increasing word).*

Proof. One can show that each L-hook or F-hook must be factored out step by step, from left to right, by applying the algorithm of rix-factorization. $\square$

For each permutation $\sigma \in \mathfrak{S}_n$ we define its *standard cycle form* (SCF) as the product of ordered cycles satisfying the following conditions:

- each cycle $(a \sigma(a) \dots \sigma^l(a))$, with $\sigma^{l+1}(a) = a$, is written with its largest element first $a \geq \sigma^k(a)$ for all $k \geq 1$;
- the cycles of length ≥ 2 are arranged in decreasing order of their largest element;
- the cycles of length 1 are arranged in increasing order of their elements and after the cycles of length ≥ 2 .

For example, if $\sigma = 129753468$, then $\text{SCF}(\sigma) = (9\ 8\ 6\ 3)(7\ 4)(1)(2)(5)$.

Let $\text{FIX}(\sigma)$ be the set of all fixed points of $\sigma \in \mathfrak{S}_n$ and introduce the set

$$\mathcal{R}_{n,k}^0 := \{\sigma \in \mathfrak{S}_n : \text{rix}(\sigma) = 0, \text{dd}(\sigma) = 1, \text{des}(\sigma) = k\}. \quad (3.2)$$

Proposition 11. *There is a bijection $\Phi : \mathfrak{S}_n \rightarrow \mathfrak{S}_n$ satisfying*

$$\text{des}(\sigma) = \text{exc}(\Phi(\sigma)) \quad \text{and} \quad \text{RIX}(\sigma) = \text{FIX}(\Phi(\sigma))$$

for each $\sigma \in \mathfrak{S}_n$. Moreover, the restriction of Φ on $\mathcal{R}_{n,k}^0$ is a bijection from $\mathcal{R}_{n,k}^0$ to $\mathcal{E}_{n,k}$.

Proof. Starting from $\sigma \in \mathfrak{S}_n$ with rix-factorization $\alpha_1 \alpha_2 \dots \alpha_i \beta$. Deleting all the rixed points of σ we get

$$\sigma' = \alpha_1 \alpha_2 \dots \alpha_i \beta',$$

where β' is the word (may be empty) obtained from β by removing all the rixed points. Clearly, β' is a F-hook. To any L-hook or F-hook $\alpha = a_1 a_2 \dots a_l$, we associate the cycle

$$\tilde{\alpha} = \begin{cases} (a_l, a_{l-1}, \dots, a_1), & \text{if } \alpha \text{ is a L-hook;} \\ (a_1, a_l, a_{l-1}, \dots, a_2), & \text{if } \alpha \text{ is a F-hook.} \end{cases}$$

Now, we define the SCF of $\Phi(\sigma)$ by

$$\Phi(\sigma) := \tilde{\alpha}_1 \tilde{\alpha}_2 \dots \tilde{\alpha}_i \tilde{\beta}'(\sigma_k)(\sigma_{k+1}) \dots (\sigma_n),$$

where $\text{RIX}(\sigma) = \{\sigma_k, \sigma_{k+1}, \dots, \sigma_n\}$. By Proposition 9, the mapping Φ is well-defined. For example, if

$$\sigma = 7\ 6\ 9\ 1\ 8\ 4\ 2\ 3\ 5\ 10 = 7\ 6\ 9|1\ 8|4\ 2\ 3\ 5\ 10,$$

then $\sigma' = 769|18|423$ and thus

$$\Phi(\sigma) = (9, 6, 7)(8, 1)(4, 3, 2)(5)(10) = 84235791610.$$

Clearly, $\text{des}(\sigma) = 4 = \text{exc}(\Phi(\sigma))$ and $\text{RIX}(\sigma) = \{5, 10\} = \text{FIX}(\Phi(\sigma))$.

To show that Φ is a bijection, we define explicitly its inverse. Given a permutation $\sigma \in \mathfrak{S}_n$, we write σ in standard cycle form as

$$\sigma = C_1 C_2 \cdots C_i O_1 O_2 \cdots O_j,$$

where the C 's are cycles of length ≥ 2 and $O_k = (o_k)$ is one point cycle for $k = 1, 2, \dots, j$. For each cycle $C = (c_1, c_2, \dots, c_l)$ of length $l \geq 2$, we define the two words (L-hook or F-hook)

$$\tilde{C} := c_l c_{l-1} \cdots c_1 \quad \text{and} \quad C' := c_1 c_l c_{l-1} \cdots c_2.$$

Then $\Phi^{-1}(\sigma)$ (viewed as a word) is defined as

$$\Phi^{-1}(\sigma) = \begin{cases} \tilde{C}_1 \tilde{C}_2 \cdots C'_i o_1 \cdots o_j, & \text{if } o_1 > \text{the largest element of } C_i; \\ \tilde{C}_1 \tilde{C}_2 \cdots \tilde{C}_i o_1 \cdots o_j, & \text{otherwise.} \end{cases}$$

For example, $\Phi^{-1}((9, 6, 7)(8, 1)(4, 3, 2)(5)(10)) = 76918423510$.

It is straightforward to check the desired properties of Φ . $\square$

Remark 2. *In view of (1.11), the two triples $(\text{exc}, \text{fix}, \text{maj})$ and $(\text{des}, \text{rix}, \text{aid})$ are equidistributed on $\mathfrak{S}_n$, where $\text{aid}(\sigma) = \text{ai}(\sigma) + \text{des}(\sigma)$. Note that (FIX, maj) and (RIX, aid) are not equidistributed on $\mathfrak{S}_3$.*

It is easy to construct a bijection between $\mathcal{R}_{n,k}^0$ and $\tilde{\mathfrak{D}}_{n,k}$. Define the mapping

$$f : \mathcal{R}_{n,k}^0 \rightarrow \tilde{\mathfrak{D}}_{n,k} \quad \text{by} \quad f(\sigma) = \varphi'_x(\sigma), \quad (3.3)$$

where σ is any permutation in $\mathcal{R}_{n,k}^0$ and $x = \beta_1(\sigma)$.

Corollary 12. *The mapping $f : \mathcal{R}_{n,k}^0 \rightarrow \tilde{\mathfrak{D}}_{n,k}$ is a bijection.*

Proof. For each $\sigma \in \mathcal{R}_{n,k}^0$, by Proposition 9 the letter $\beta_1(\sigma)$ is the only double descent of σ , which becomes a double ascent of $f(\sigma)$. As $\text{rix}(\sigma) = 0$, the last letter of $f(\sigma)$ is $\beta_1(\sigma)$. Thus $f(\sigma)$ is a permutation in $\tilde{\mathfrak{D}}_{n,k}$ and f is well-defined. To show that f is a bijection we define its inverse f^{-1} as the word

$$f^{-1}(\sigma) := \varphi'_y(\sigma),$$

where σ is any permutation in $\tilde{\mathfrak{D}}_{n,k}$ and y is the last letter of σ . All we need to check is that now $f^{-1}(\sigma)$ is a permutation in $\mathcal{R}_{n,k}^0$. Clearly, the letter y becomes the only double descent in $f^{-1}(\sigma)$ and $\text{des}(f^{-1}(\sigma)) = k$. Moreover, we have $y = \beta_1(f^{-1}(\sigma))$ by the algorithm of the rix -factorization, and so $\text{rix}(f^{-1}(\sigma)) = 0$. Therefore, the word $f^{-1}(\sigma)$ is a permutation in $\mathcal{R}_{n,k}^0$. $\square$

Remark 3. In view of Proposition 11, the composition $\Phi \circ f^{-1}$ is a bijection between $\tilde{\mathfrak{D}}_{n,k}$ and $\mathcal{E}_{n,k}$. This bijection is illustrated in Table 1.

$\tilde{\mathfrak{D}}_{4,2}$	1324	1423	2314	2413	3412
$\mathcal{R}_{4,2}^0$	4132	1432	4213	2431	3421
$\mathcal{E}_{4,2}$	4312	4321	2413	3412	2143

TABLE 1. The bijection $\Phi \circ f^{-1} : \tilde{\mathfrak{D}}_{4,2} \xrightarrow{f^{-1}} \mathcal{R}_{4,2} \xrightarrow{\Phi} \mathcal{E}_{4,2}$.

4. PROOF OF THEOREM 5

For any $x \in [n]$, we introduce a new action $\varphi_x'' : \mathfrak{S}_n \rightarrow \mathfrak{S}_n$ as

$$\varphi_x''(\sigma) = \begin{cases} \sigma, & \text{if } x \in \{\beta_1(\sigma)\} \cup \text{RIX}(\sigma); \\ \varphi_x'(\sigma), & \text{otherwise.} \end{cases}$$

Since φ_x' 's are involutions and that they commute, so do φ_x'' 's. Thus, for any subset $S \subseteq [n]$ we can define the mapping $\varphi_S'' : \mathfrak{S}_n \rightarrow \mathfrak{S}_n$ by

$$\varphi_S''(\sigma) = \prod_{x \in S} \varphi_x''(\sigma).$$

This is a new $\mathbb{Z}_2^n$ -action on $\mathfrak{S}_n$ via the functions φ_S'' , $S \subseteq [n]$, that we called *restricted MFS-action*. An important property of the action φ_x'' is that it preserves the rix-factorization type of permutations, namely,

Lemma 13. Let $\sigma \in \mathfrak{S}_n$ be a permutation with rix-factorization $\alpha_1\alpha_2\cdots\alpha_i\alpha_{i+1}$ with $\alpha_{i+1} = \beta$. For any $x \in [n]$, if x is a letter of α_k ($1 \leq k \leq i+1$), then the rix-factorization of $\varphi_x''(\sigma)$ is $\alpha_1\cdots\alpha'_k\cdots\alpha_i\alpha_{i+1}$, where α'_k is a rearrangement of α_k . Moreover,

$$\beta_1(\varphi_x''(\sigma)) = \beta_1(\sigma) \quad \text{and} \quad \text{RIX}(\varphi_x''(\sigma)) = \text{RIX}(\sigma). \quad (4.1)$$

Proof. For $x \in [n]$ we apply φ_x'' to σ and distinguish two cases.

- x is a letter of α_k ($1 \leq k \leq i$). Let x_k be the last letter of α_k , which is also the greatest letter in α_k . If $x = x_k$, then x is a peak of σ by Proposition 9, so $\varphi_x''(\sigma) = \sigma$. If $x \neq x_k$, then $x < x_k < x_{k-1}$ by Proposition 9, where $x_0 = +\infty$. Therefore, $\varphi_x''(\sigma) = \alpha_1\cdots\alpha_{k-1}\varphi_x'(\alpha_k)\alpha_{k+1}\cdots\alpha_i\beta$ with x_k as the last letter of $\varphi_x'(\alpha_k)$.
- x is a letter of $\alpha_{i+1} = \beta$. If $x \in \{\beta_1(\sigma)\} \cup \text{RIX}(\sigma)$, then $\varphi_x''(\sigma) = \sigma$; otherwise, $x < \beta_1(\sigma)$. Therefore, $\varphi_x''(\sigma) = \alpha_1\cdots\alpha_i\varphi_x'(\beta)$ with $\beta_1(\sigma)$ as the first letter of $\varphi_x'(\beta)$.

In both cases, the result follows from the rix-factorization. $\square$

Let $\mathcal{R}_n$ be the set of permutations in $\mathfrak{S}_n$ without rixed points.

Lemma 14. *Let “st” be a statistic invariant under the restricted MFS-action. Then,*

$$\sum_{\sigma \in \mathcal{R}_n} q^{\text{st}(\sigma)} t^{\text{des}(\sigma)} = \sum_{k=1}^{\lfloor n/2 \rfloor} \left(\sum_{\sigma \in \mathcal{R}_{n,k}^0} q^{\text{st}(\sigma)} \right) t^k (1+t)^{n-2k}.$$

Proof. First we note that the restricted MFS-action is stable on $\mathcal{R}_n$, that is, $\sigma \in \mathcal{R}_n$ implies $\varphi_x''(\sigma) \in \mathcal{R}_n$ for any $x \in [n]$. This follows from the property of φ_x'' in (4.1).

For a permutation $\sigma \in \mathcal{R}_n$ let $\widetilde{\text{Orb}}(\sigma) = \{g(\sigma) : g \in \mathbb{Z}_2^n\}$ be the orbit of σ in $\mathcal{R}_n$ under the restricted MFS-action. By Proposition 9, the letter $\beta_1(\sigma)$ is a double descent of σ . If $x \in [n]$ is a double descent of σ different from $\beta_1(\sigma)$, then x becomes a double ascent of $\varphi_x''(\sigma)$ and $\text{des}(\sigma) = \text{des}(\varphi_x''(\sigma)) + 1$. Therefore, there is a unique element $\tilde{\sigma} \in \widetilde{\text{Orb}}(\sigma)$ with one double descent $\beta_1(\tilde{\sigma})$ such that

$$\sum_{\pi \in \widetilde{\text{Orb}}(\sigma)} q^{\text{st}(\pi)} t^{\text{des}(\pi)} = q^{\text{st}(\tilde{\sigma})} t^{\text{des}(\tilde{\sigma})} (1+t)^{\text{da}(\tilde{\sigma})}.$$

Since, for any $\sigma \in \mathfrak{S}_n$, we have $\text{valley}(\sigma) = \text{peak}(\sigma) + 1$ and $\text{dd}(\sigma) + \text{da}(\sigma) + 2\text{peak}(\sigma) = n - 1$, the result follows then from the fact that $\text{dd}(\tilde{\sigma}) = 1$ and $\text{peak}(\tilde{\sigma}) = \text{des}(\tilde{\sigma}) - 1$. $\square$

It follows from Eq. (1.11) and Lemmas 7 and 14 that

$$A_n(t, 0, q) = \sum_{\sigma \in \mathcal{R}_n} t^{\text{des}(\sigma)} q^{\text{ai}(\sigma)} = \sum_{k=1}^{\lfloor n/2 \rfloor} \left(\sum_{\sigma \in \mathcal{R}_{n,k}^0} q^{\text{ai}(\sigma)} \right) t^k (1+t)^{n-2k}.$$

Applying the bijection $f : \mathcal{R}_{n,k}^0 \rightarrow \tilde{\mathfrak{D}}_{n,k}$ defined in (3.3), we get

$$A_n(t, 0, q) = \sum_{k=1}^{\lfloor n/2 \rfloor} \left(\sum_{\sigma \in \tilde{\mathfrak{D}}_{n,k}} q^{\text{ai}(\sigma)} \right) t^k (1+t)^{n-2k},$$

which is equivalent to expansion (1.10) in view of Lemma 8.

5. GENERATING FUNCTIONS OF γ -COEFFICIENTS

In this section, we derive the recurrences and generating functions of the two γ -coefficients in (1.1) and (1.2) from their combinatorial interpretations (1.9) and (1.10), namely,

$$\gamma_{n,k}(q) = \sum_{\sigma \in \mathfrak{D}_{n,k}} q^{\text{inv}(\sigma)} \quad \text{and} \quad \tilde{\gamma}_{n,k}(q) = \sum_{\sigma \in \tilde{\mathfrak{D}}_{n,k}} q^{\text{inv}(\sigma)}. \quad (5.1)$$

Let $\mathfrak{D}_n = \{\sigma \in \mathfrak{S}_n : \text{dd}(\sigma) = 0\}$ and $\tilde{\mathfrak{D}}_n = \{\sigma \in \mathfrak{D}_n : \sigma_{n-1} < \sigma_n\}$. Then

$$\Gamma_n(y, q) := \sum_{k=0}^{\lfloor (n-1)/2 \rfloor} \gamma_{n,k}(q) y^k = \sum_{\sigma \in \mathfrak{D}_n} y^{\text{des}(\sigma)} q^{\text{inv}(\sigma)}; \quad (5.2)$$

$$\tilde{\Gamma}_n(y, q) := \sum_{k=1}^{\lfloor n/2 \rfloor} \tilde{\gamma}_{n,k}(q) y^k = \sum_{\sigma \in \tilde{\mathfrak{D}}_n} y^{\text{des}(\sigma)+1} q^{\text{inv}(\sigma)}. \quad (5.3)$$

The first values of $\Gamma_n(y, q)$ and $\tilde{\Gamma}_n(y, q)$ for $1 \leq n \leq 5$ are:

$$\begin{aligned} \Gamma_1(y, q) &= \Gamma_2(y, q) = 1, & \Gamma_3(y, q) &= 1 + y(q + q^2), \\ \Gamma_4(y, q) &= 1 + y(q + q^2)(2 + q + q^2), \\ \Gamma_5(y, q) &= 1 + y(3q + 5q^2 + 5q^3 + 5q^4 + 2q^5 + 2q^6) \\ &\quad + y^2(q + q^3)(1 + q + q^2 + q^3)(q + q^2); \end{aligned}$$

and

$$\begin{aligned} \tilde{\Gamma}_1(y, q) &= 0, & \tilde{\Gamma}_2(y, q) &= \tilde{\Gamma}_3(y, q) = y, \\ \tilde{\Gamma}_4(y, q) &= y + y^2(q + 2q^2 + q^3 + q^4), \\ \tilde{\Gamma}_5(y, q) &= y + y^2(2q + 4q^2 + 4q^3 + 4q^4 + 2q^5 + 2q^6). \end{aligned}$$

For $n \geq 1$ let $(q; q)_n := \prod_{i=1}^n (1 - q^i)$ be the shifted q -factorial and $(q; q)_0 = 1$. The q -exponential function and q -binomial coefficients are then defined by

$$e(z; q) := \sum_{n \geq 0} \frac{z^n}{(q; q)_n} \quad \text{and} \quad \begin{bmatrix} n \\ k \end{bmatrix}_q := \frac{(q; q)_n}{(q; q)_k (q; q)_{n-k}}.$$

The following generating function formula is due to Shareshian and Wachs [15, 16]:

$$1 + \sum_{n \geq 1} A_n(t, r, q) \frac{z^n}{(q; q)_n} = \frac{(1-t)e(rz; q)}{e(tz; q) - te(z; q)}. \quad (5.4)$$

Proposition 15. *The exponential generating functions for $\Gamma_n(y, q)$ and $\tilde{\Gamma}_n(y, q)$ are*

$$\sum_{n \geq 0} \Gamma_n(y, q) \frac{z^n}{(q; q)_n} = \frac{e(z/(1+t); q) - te(tz/(1+t); q)}{e(tz/(1+t); q) - te(z/(1+t); q)}, \quad (5.5)$$

$$\sum_{n \geq 0} \tilde{\Gamma}_n(y, q) \frac{z^n}{(q; q)_n} = \frac{1-t}{e(tz/(1+t); q) - te(z/(1+t); q)}, \quad (5.6)$$

where $\Gamma_0(y, q) = \tilde{\Gamma}_0(y, q) = 1$ and $y = t/(1+t)^2$.

Proof. Clearly Theorems 4 and 5 are equivalent to, for $n \geq 1$,

$$A_n(t, 1, q) = (1 + t)^{n-1} \Gamma_n(y, q), \quad (5.7)$$

$$A_n(t, 0, q) = (1 + t)^n \tilde{\Gamma}_n(y, q), \quad (5.8)$$

where $y = t/(1 + t)^2$. Substituting (5.7) and (5.8) into (5.4) yields (5.5) and (5.6). $\square$

Recall the well-known combinatorial interpretation of the q -binomial coefficients [19, Prop. 1.3.17]

$$\begin{bmatrix} n \\ k \end{bmatrix}_q = \sum_{(\mathcal{A}, \mathcal{B})} q^{\text{inv}(\mathcal{A}, \mathcal{B})}, \quad (5.9)$$

where the sum is over all ordered set partitions $(\mathcal{A}, \mathcal{B})$ of $[n]$ such that $|\mathcal{A}| = k$ and

$$\text{inv}(\mathcal{A}, \mathcal{B}) := \{(i, j) \in \mathcal{A} \times \mathcal{B} : i > j\}.$$

Proposition 16. *The polynomials $\Gamma_n(y, q)$ and $\tilde{\Gamma}_n(y, q)$ satisfy the following recurrences:*

$$\Gamma_{n+1}(y, q) = \Gamma_n(y, q) + y \sum_{i=1}^{n-1} q^i \begin{bmatrix} n \\ i \end{bmatrix}_q \Gamma_i(y, q) \Gamma_{n-i}(y, q); \quad (5.10)$$

$$\tilde{\Gamma}_{n+1}(y, q) = \Gamma_n(y, q) + y \sum_{i=2}^{n-1} q^i \begin{bmatrix} n \\ i \end{bmatrix}_q \tilde{\Gamma}_i(y, q) \Gamma_{n-i}(y, q); \quad (5.11)$$

for $n \geq 1$ and the initial conditions $\Gamma_0(t, q) = \Gamma_1(t, q) = \tilde{\Gamma}_0(t, q) = 1$ and $\tilde{\Gamma}_1(t, q) = 0$.

Proof. For $2 \leq j \leq n$ let $\mathfrak{D}_n^{(j)} := \{\sigma \in \mathfrak{D}_n : \sigma_j = n\}$ and define

$$\Gamma_n^{(j)}(y, q) := \sum_{\sigma \in \mathfrak{D}_n^{(j)}} y^{\text{des}(\sigma)} q^{\text{inv}(\sigma)}.$$

Clearly, for $n \geq 1$

$$\Gamma_{n+1}(y, q) = \sum_{j=2}^{n+1} \Gamma_{n+1}^{(j)}(y, q) = \Gamma_n(y, q) + \sum_{j=2}^n \Gamma_n^{(j)}(y, q). \quad (5.12)$$

For any finite ordered set X let $\binom{X}{m}$ denote the set of the m -element subsets of X and $\mathfrak{D}_X$ the set of permutations of X without double descents. Also let $\mathcal{T}(n, j)$ be the set of all triples (S, σ_L, σ_R) such that $S \in \binom{[n]}{j}$ and $\sigma_L \in \mathfrak{D}_S$, $\sigma_R \in \mathfrak{D}_{[n] \setminus S}$. For $2 \leq j \leq n-1$ and $\sigma \in \mathfrak{D}_n^{(j)}$, define the mapping $\sigma \mapsto (S, \sigma_L, \sigma_R)$ by

- $S = \{\sigma_i : 1 \leq i \leq j-1\}$;
- $\sigma_L = \sigma_1 \sigma_2 \cdots \sigma_{j-1}$ and $\sigma_R = \sigma_{j+1} \sigma_{j+2} \cdots \sigma_n$.

It is not hard to see that this mapping is a bijection between $\mathfrak{D}_n^{(j)}$ and $\mathcal{T}(n-1, j-1)$ and satisfies

$$\text{des}(\sigma) = \text{des}(\sigma_L) + \text{des}(\sigma_R) + 1$$

and

$$\text{inv}(\sigma) = \text{inv}(\sigma_L) + \text{inv}(\sigma_R) + \text{inv}(S, [n-1] \setminus S) + n - j.$$

Thus, for $2 \leq j \leq n$ we have

$$\begin{aligned} \Gamma_{n+1}^{(j)}(y, q) &= \sum_{\sigma \in \mathfrak{D}_{n+1}^{(j)}} y^{\text{des}(\sigma)} q^{\text{inv}(\sigma)} \\ &= yq^{n+1-j} \sum_{(S, \sigma_L, \sigma_R) \in \mathcal{T}(n, j-1)} q^{\text{inv}(S, [n] \setminus S)} q^{\text{inv}(\sigma_L)} y^{\text{des}(\sigma_L)} q^{\text{inv}(\sigma_R)} y^{\text{des}(\sigma_R)} \\ &= yq^{n+1-j} \sum_{S \in \binom{[n]}{j-1}} q^{\text{inv}(S, [n] \setminus S)} \sum_{\sigma_L \in \mathfrak{D}_S} q^{\text{inv}(\sigma_L)} y^{\text{des}(\sigma_L)} \sum_{\sigma_R \in \mathfrak{D}_{[n] \setminus S}} q^{\text{inv}(\sigma_R)} y^{\text{des}(\sigma_R)} \\ &= yq^{n+1-j} \left[\begin{matrix} n \\ j-1 \end{matrix} \right]_q \Gamma_{j-1}(y, q) \Gamma_{n+1-j}(y, q), \end{aligned} \quad (5.13)$$

where we apply (5.9) to the last equality. Substituting (5.13) into (5.12) we obtain (5.10). Similarly we can prove (5.11), the details are left to the interested reader. $\square$

Remark 4. We can also prove Proposition 16 from the following known recurrence of $A_n(t, r, q)$ ($n \geq 1$), which was proved in [10] by applying the q -differential operator to both sides of (5.4),

$$A_{n+1}(t, r, q) = rA_n(t, r, q) + t \sum_{j=0}^{n-1} \left[\begin{matrix} n \\ j \end{matrix} \right]_q q^j A_j(t, r, q) A_{n-j}(t, 1, q) \quad (5.14)$$

with initial condition $A_0(t, r, q) = 1, A_1(t, r, q) = r$. Substituting (5.7) and (5.8) into (5.14) we also obtain the above recurrences for γ -coefficients.

6. CONCLUDING REMARKS

Using the combinatorics developed in Section 3, we can also give a combinatorial proof of Theorem 2, of which a similar proof for the special $\beta = 1$ case was given in [1, 20].

Proof of Theorem 2. Define the statistic “lyc” by $\text{lyc}(\sigma) := \text{cyc}(\Phi(\sigma))$ for $\sigma \in \mathfrak{S}_n$. Since the function $\varphi_x'' : \mathfrak{S}_n \rightarrow \mathfrak{S}_n$ preserves the rix-factorization type of permutations for any $x \in [n]$, the statistic “lyc” is invariant under the restricted MFS-action. Thus by Lemma 14, we have

$$\sum_{\sigma \in \mathcal{R}_n} \beta^{\text{lyc}(\sigma)} t^{\text{des}(\sigma)} = \sum_{j=1}^{\lfloor n/2 \rfloor} \left(\sum_{\sigma \in \mathcal{R}_{n,j}^0} \beta^{\text{lyc}(\sigma)} \right) t^j (1+t)^{n-2j}.$$

By Proposition 11, applying the bijection $\Phi : \mathfrak{S}_n \rightarrow \mathfrak{S}_n$ to both sides of the above expansion we get (1.7). $\square$

We can further extend (1.7) and (1.10) to permutations with a given number of fixed points. For $0 \leq j \leq n$ define the set

$$\mathfrak{S}_{n,j} := \{\sigma \in \mathfrak{S}_n : \text{fix}(\sigma) = j\}.$$

Then, it is easy to see that

$$\sum_{\sigma \in \mathfrak{S}_{n,j}} \beta^{\text{cyc}(\sigma)} t^{\text{exc}(\sigma)} = \binom{n}{j} \beta^j \sum_{\sigma \in \mathfrak{S}_{n-j,0}} \beta^{\text{cyc}(\sigma)} t^{\text{exc}(\sigma)}. \quad (6.1)$$

Also, it is known [16, Eq.(4.3)] and easy to deduce from (5.4) that

$$\sum_{\sigma \in \mathfrak{S}_{n,j}} t^{\text{exc}(\sigma)} q^{\text{maj}(\sigma) - \text{exc}(\sigma)} = \left[\begin{matrix} n \\ j \end{matrix} \right]_q A_{n-j}(t, 0, q). \quad (6.2)$$

We derive from Theorem 2 and Theorem 5 the following result.

Proposition 17. *For $1 \leq j \leq n$ we have*

$$\sum_{\sigma \in \mathfrak{S}_{n,j}} \beta^{\text{cyc}(\sigma)} t^{\text{exc}(\sigma)} = \sum_{k=1}^{\lfloor n/2 \rfloor} \binom{n}{j} \left(\sum_{\sigma \in \mathcal{E}_{n-j,k}} \beta^{\text{cyc}(\sigma)+j} \right) t^k (1+t)^{n-j-2k}, \quad (6.3)$$

and

$$\sum_{\sigma \in \mathfrak{S}_{n,j}} t^{\text{exc}(\sigma)} q^{\text{maj}(\sigma) - \text{exc}(\sigma)} = \sum_{k=1}^{\lfloor (n-j)/2 \rfloor} \left[\begin{matrix} n \\ j \end{matrix} \right]_q \left(\sum_{\sigma \in \tilde{\mathfrak{D}}_{n-j,k}} q^{\text{inv}(\sigma)} \right) t^k (1+t)^{n-j-2k}. \quad (6.4)$$

Shareshian and Wachs [16, Remark 5.5] also proved the γ -positivity of the left-hand side of (6.4) and further generalized (1.2) to

$$\sum_{\sigma \in \mathfrak{S}_{n,0}} t^{\text{exc}(\sigma)} p^{\text{des}(\sigma)} q^{\text{maj}(\sigma) - \text{exc}(\sigma)} = \sum_{k=0}^{\lfloor (n-1)/2 \rfloor} \tilde{\gamma}_{n,k}(p, q) t^k (1+t)^{n-1-2k}, \quad (6.5)$$

where $\tilde{\gamma}_{n,k}(p, q)$ are polynomials in $\mathbb{N}[p, q]$. Obviously we have $\tilde{\gamma}_{n,k}(1, q) = \tilde{\gamma}_{n,k}(q)$. It would be interesting to find a combinatorial interpretation for the polynomials $\tilde{\gamma}_{n,k}(p, q)$ in the light of (1.10).

Acknowledgement. The first author was supported by the China Scholarship Council to study abroad from 2010 to 2014 and would like to thank Institut Camille Jordan and Université Claude Bernard Lyon 1, where most of this work was done, for hosting him.

REFERENCES

- [1] C.A. Athanasiadis and C. Savvidou, The local h -vector of the cluster subdivision of a simplex, *Sem. Lothar. Combin.*, **B66c** (2012), 21pp.
- [2] P. Brändén, Actions on permutations and unimodality of descent polynomials, *European J. Combin.*, **29** (2008), 514–531.
- [3] A. Burstein, On the distribution of some Euler–Mahonian statistics, [arXiv:1402.3619](#).
- [4] D. Foata and G.-N. Han, The q -Tangent and q -Secant numbers via basic Eulerian polynomials, *Proc. Amer. Math. Soc.*, **138** (2009), 385–393.
- [5] D. Foata and M.-P. Schützenberger, *Théorie géométrique des polynômes eulériens*, Lecture Notes in Mathematics, Vol. 138, Springer-Verlag, Berlin, 1970.
- [6] D. Foata and M.-P. Schützenberger, Major index and inversion number of permutations, *Math. Machr.*, **83** (1978), 143–159.
- [7] D. Foata and V. Strehl, Rearrangements of the symmetric group and enumerative properties of the tangent and secant numbers, *Math. Z.*, **137** (1974), 257–264.
- [8] I.M. Gessel, A coloring problem, *Amer. Math. monthly*, **98** (1991), 530–533.
- [9] G.-N. Han, F. Jouhet and J. Zeng, Two new triangles of q -integers via q -Eulerian polynomials of type A and B, *Ramanujan J.*, **31** (2013), 115–127.
- [10] Z. Lin, On some generalized q -Eulerian polynomials, *Electron. J. Combin.*, **20**(1) (2013), #P55.
- [11] S. Linusson, J. Shareshian, M.L. Wachs, Rees products and lexicographic shellability, *J. Comb.*, **3** (2012), 243–276.
- [12] E. Nevo, T.K. Petersen and B.E. Tenner, The γ -vector of a barycentric subdivision. *J. Combin. Theory Ser. A* 118 (2011), no. 4, 1364–1380.
- [13] A. Postnikov, V. Reiner and L. Williams, Faces of generalized permutohedra, *Doc. Math.* 13 (2008) 207–273.
- [14] L. Shapiro, W.J. Woan, S. Getu: Runs, slides and moments. *SIAM J. Algebraic Discrete Methods* 4 (1983), 459–466.
- [15] J. Shareshian and M.L. Wachs, q -Eulerian polynomials: Excedance Number and Major Index, *Electron. Res. Announc. Amer. Math. Soc.* **13** (2007), 33–45.
- [16] J. Shareshian and M.L. Wachs, Eulerian quasisymmetric functions, *Adv. in Math.*, **225** (2011), 2921–2966.
- [17] H. Shin and J. Zeng, The symmetric and unimodal expansion of eulerian polynomials via continued fractions, *European J. Combin.*, **33** (2012), 111–127.
- [18] H. Shin and J. Zeng, Symmetric unimodal expansions of excedances in colored permutations, [arXiv:1411.6184](#).
- [19] R.P. Stanley, *Enumerative combinatorics*, Vol. 1, Cambridge University Press, Cambridge, 1997.
- [20] H. Sun and Y. Wang, A group action on derangements, *Electron. J. Combin.*, **21**(1) (2014), #P1.67.

(Zhicong Lin) SCHOOL OF SCIENCES, JIMEI UNIVERSITY, XIAMEN 361021, P.R. CHINA
E-mail address: lin@math.univ-lyon1.fr

(Jiang Zeng) INSTITUT CAMILLE JORDAN, UNIVERSITÉ CLAUDE BERNARD LYON 1, FRANCE
E-mail address: zeng@math.univ-lyon1.fr